**On the Maximum Sequence in a Critical Branching Process**

K. B. Athreya

*The Annals of Probability*, Vol. 16, No. 2 (Apr., 1988), 502-507.

Stable URL:
http://links.jstor.org/sici?sici=0091-1798%28198804%2916%3A2%3C502%3AOTMSIA%3E2.0.CO%3B2-O

*The Annals of Probability* is currently published by Institute of Mathematical Statistics.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/about/terms.html. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/journals/ims.html.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

http://www.jstor.org/
Sat Mar 12 05:05:47 2005

*The Annals of Probability*
1988, Vol. 16, No. 2, 502–507

# ON THE MAXIMUM SEQUENCE IN A CRITICAL BRANCHING PROCESS[1]

BY K. B. ATHREYA

*Iowa State University, Indian Statistical Institute and the Mathematical Sciences Research Institute, Berkeley*

If $\{Z_n\}_0^\infty$ is a critical branching process such that $E_1 Z_1^2 < \infty$, then $(\log n)^{-1} E_i M_n \to i$, where $E_i$ refers to starting with $Z_0 = i$ and $M_n = \max_{0 \le j \le n} Z_j$. This improves the earlier results of Weiner [9] and Pakes [7].

**1. Introduction.** Let $\{Z_n\}_0^\infty$ be a critical branching process. (See [1] for a complete definition and basic results.) Let $M_n = \max_{0 \le j \le n} Z_j$. This paper studies the growth rate of $EM_n$.

Here is a quick argument to show that $EM_n \to \infty$. By monotone convergence theorem $EM_n \to EM$, where $M = \max_{j \ge 0} Z_j$. Clearly,

$$E\{Z_n \colon Z_n > 0\} \le E(M \colon Z_n > 0),$$

and if $EM < \infty$, then this goes to zero since $P(Z_n > 0) \to 0$. On the other hand, $E\{Z_n \colon Z_n > 0\} = EZ_n = EZ_0$ for all $n$.

We establish the following:

THEOREM 1. *If $E_1 Z_1^2 < \infty$, then $(\log n)^{-1} E_i M_n \to i$, where $E_i$ refers to starting with $Z_0 = i$.*

This is an improvement over the results of Weiner [9] and Pakes [7]. The former showed that if there exists a finite $K$ such that $P_1(Z_1 < K) = 1$, then there exists a finite positive constant $\alpha_K$ such that $EM_n \ge \alpha_K \log n$ with $\alpha_K \to 0$ as $K \to \infty$, and that if $E_1 Z_1^2 < \infty$, then there exists a $\beta$ such that $EM_n \le \beta \log n$. Pakes [7] improved Weiner's upper bound and showed that

$$\limsup_n (EM_n)(\log n)^{-1} \le 1, \quad \text{if } E_1(Z_1^2) < \infty.$$

Here is an outline of the proof of Theorem 1.

*Step* 1. Pakes [7] has shown that for any nonnegative martingale $\{Z_n\}$ such that $E(Z_n(\log Z_n)^+) \to \infty$, Doob's maximal inequality ([2], page 263) can be improved to get

$$\limsup_n EM_n\big(EZ_n(\log Z_n)^+\big)^{-1} \le 1.$$

Received April 1986; revised April 1987.

[1]Research supported in part by NSF Grant DMS-85-02311, by a visiting professorship at the Indian Statistical Institute, Bangalore, and by a visiting membership at the Mathematical Sciences Research Institute, Berkeley.

*AMS* 1980 *subject classifications.* 60J80, 60K99.

*Key words and phrases.* Branching process, critical, maximum.

Since a critical branching process $\{Z_n\}$ is a nonnegative martingale for which $EZ_n(\log Z_n)^+$ does go to $\infty$ (see Proposition 1) this result is available to us without any additional moment conditions other than $E_1Z_1 = 1$.

*Step* 2. Using the conditional limit law of Yaglom (see [1], page 20) we establish (see Proposition 1) that if $E_1Z_1^2 < \infty$, then

$$\lim_n \left(E_iZ_n(\log Z_n)^+ - i\log n\right) = \mu_i, \qquad 0 < \mu_i < \infty,$$

exists for each $i \geq 1$.

*Step* 3. Using a stopping time argument we establish a bound for the lim inf (see Proposition 4).

If $E_1Z_1^2 < \infty$, then

$$\liminf_n E_iM_n(\log n)^{-1} \geq i.$$

Clearly, the above three steps imply Theorem 1.

## 2. Proofs.

PROPOSITION 1. *Let $\{Z_n\}_0^\infty$ be a critical branching process with offspring distribution $\{p_j\}$. Assume $p_1 \neq 1$. Then* (i) *$E_iZ_n\log Z_n \to \infty$ for each $i \geq 1$, and* (ii) *if, in addition, $\sum j^2p_j < \infty$, then for $i \geq 0$, $\lim_n(E_iZ_n\log Z_n - i\log n) = i\mu$ exists, where $0 < \mu < \infty$.*

PROOF. (i) It is enough to consider $i = 1$,

$$\begin{aligned} EZ_n\log Z_n &= E(Z_n\log Z_n|Z_n > 0)P(Z_n > 0) \\ &= E\left(a_n^{-1}Z_n\log\left(a_n^{-1}Z_n\right)|Z_n > 0\right) + \log a_n, \end{aligned}$$

if we choose $a_n = (P(Z_n > 0))^{-1}$.

Thus, $EZ_n\log Z_n - \log a_n \geq E(a_n^{-1}Z_n\log(a_n^{-1}Z_n)\colon\ 0 < Z_n < a_n|Z_n > 0)$. But since $\sup_{0<x<1}|x\log x| = C < \infty$, we get

$$\liminf(EZ_n\log Z_n - \log a_n) > -C,$$

proving (i) as $a_n \to \infty$.

(ii) Since $E_iZ_n = i$,

$$\begin{aligned} E_iZ_n\log Z_n - i\log n &= E_i(Z_n\log Z_n - Z_n\log n) \\ &= E_i\left(n^{-1}Z_n\log\left(n^{-1}Z_n\right)|Z_n > 0\right)nP_i(Z_n > 0). \end{aligned}$$

From (2) we see that $nP_i(Z_n > 0) \to i\lambda$. Now

$$\begin{aligned} E_i\left(n^{-2}Z_n^2|Z_n > 0\right) &= n^{-1}E_i\left(Z_n^2\colon\ Z_n > 0\right)\left(nP_i(Z_n > 0)\right)^{-1} \\ &= n^{-1}\left(in\sigma^2 + i^2\right)\left(nP_i(Z_n > 0)\right)^{-1}, \end{aligned}$$

and this converges to $2^{-1}\sigma^4$ as $n \to \infty$. Thus, $n^{-1}Z_n\log(n^{-1}Z_n)|Z_n > 0$ are uniformly integrable and (3) now implies (ii) with $\mu = \lambda\int_0^\infty(x\log x)\lambda e^{-\lambda x}\,dx$. □

PROPOSITION 2. *Let $\{S_k\}_0^\infty$ be random walk generated by the probability distribution $\{p_j\}_0^\infty$. Let $\sum jp_j = 1$ and $\sum j^2 p_j < \infty$. Let $\rho > 1$ and $\psi(k, l) = E(S_k\colon S_k > \rho l)$. Then $\sum_l l^{-1}\psi(l, l) < \infty$.*

PROOF. It is enough to consider the case $\rho = (1 + k^{-1})$ when $k$ is a positive integer. Clearly,

$$\begin{aligned} l^{-1}\psi(l, l) &= l^{-1}E(S_l\colon S_l > \rho l) \\ &= l^{-1}E(S_l - \rho l\colon S_l > \rho l) + \rho P(S_l > \rho l) \\ &= a_l + b_l, \quad \text{say.} \end{aligned}$$

Since $\sum_j j^2 p_j < \infty$, $\sum_l b_l \le \rho \sum_l P(|S_l - l| > l/k) < \infty$ by the criterion for complete convergence due to Hsu and Robbins [5].

Next,

$$\begin{aligned} kla_l &= kE(S_l - \rho l\colon S_l > \rho l) \\ &= E(S_l'\colon S_l' > 0), \end{aligned}$$

where $\{S_l'\}$ is a random walk generated by the random variable $S_1' = k(S_1 - 1) - 1$. From Spitzer [8], pages 180–181, we know that

$$\sum_l l^{-1}E(S_l'\colon S_l' > 0) = \frac{E(S_T'\colon T < \infty)}{P(T = \infty)},$$

where

$$T = \begin{cases} \inf\{n\colon n \ge 1, S_n' > 0\}, \\ \infty, \quad \text{if } S_n' \le 0 \text{ for all } n \ge 1. \end{cases}$$

Since $ES_1' = -1 < 0$, $p = P(T = \infty) > 0$. Also $E(S_T'\colon T < \infty) \le EM'$, where $M' = \sup\{S_n'\colon n \ge 0\}$. Kiefer and Wolfowitz [6] have shown that $EM' < \infty$ if $ES_1'^2 < \infty$. But $ES_1'^2 < \infty$ is the same as $\sum j^2 p_j < \infty$. Thus, $\sum_l a_l < \infty$ and we are done. □

REMARK 1. Actually, the converse to Proposition 2 is also valid. If $\sum_l l^{-1}\psi(l, \rho l) < \infty$ for some $\rho > 1$, then $\sum b_l < \infty$ implying $\sum j^2 p_j < \infty$ by a converse proved by Erdös [3] to the Hsu and Robbins result [5] on complete convergence.

PROPOSITION 3. *Let $\{Z_n\colon n = 0, 1, 2 \dots\}$ be a nonnegative martingale. Fix $n \ge 1$, $l > 1$. Let $T = T_{n,l}$ be a stopping time defined by*

$$T = \begin{cases} \min\{r\colon 1 \le r \le n, Z_r = 0 \text{ or } Z_r \ge l\}, \\ n, \quad \text{if } 0 < Z_r < l \text{ for } 1 \le r \le n. \end{cases}$$

*Then $E(Z_T\colon Z_T \ge l) \ge E(Z_n\colon Z_n \ge l)$.*

PROOF. Since $T$ is a bounded stopping time for the martingale $\{Z_n\}$, the optional sampling theorem applies so that $E(Z_T) = E(Z_n)$. The event

$(0 < Z_T < l)$ is the same as $T = n$ and $0 < Z_n < l$. Thus, $E(Z_T\colon 0 < Z_T < l) = E(Z_n\colon 0 < Z_n < l,\ T = n) \le E(Z_n\colon 0 < Z_n < l)$. Since $E(Z_T) = E(Z_n)$ the given assertion follows. □

PROPOSITION 4. *Let $\{Z_n\}$ be a critical branching process with offspring distribution $\{p_j\}$ such that $\sum j^2 p_j < \infty$. Then*

$$\liminf_n (\log n)^{-1} E_i M_n \ge i.$$

PROOF. Now let $\{Z_n\}$ be a critical branching process. It is a nonnegative martingale and so Proposition 3 applies. For fixed integers $l$ and $n$ and positive number $\rho > 1$, we get

$$\begin{aligned} E(Z_n\colon Z_n \ge l) &\le E(Z_T\colon Z_T \ge l) \\ &= E(Z_T\colon Z_T \ge l,\ Z_T \le \rho l) + E(Z_T\colon Z_T \ge l,\ Z_T > \rho l) \\ &= A_{n,l} + B_{n,l}, \quad \text{say.} \end{aligned}$$

Clearly, $A_{n,l} \le \rho l P(M_n \ge l)$, where $M_n = \max_{0 \le j \le n}\{Z_j\}$. Next

$$\begin{aligned} B_{n,l} &= \sum_{r=1}^{n} E(Z_T\colon Z_T > \rho l;\ T = r) \\ &= \sum_{r=1}^{n} E(Z_r\colon Z_r > \rho l;\ Z_1 < l, \ldots, Z_{r-1} < l) \\ &\le \sum_{r=1}^{n} E(\psi(Z_{r-1}, l);\ 0 < Z_{r-1} < l), \end{aligned}$$

where $\psi(k, l) = E(S_k\colon S_k > \rho l)$, $\{S_k\}_0^\infty$ being random walk generated by the offspring distribution $\{p_j\}$. Since $\psi(k, l)$ is increasing in $k$ for fixed $l$, we have

$$B_{n,l} \le \psi(l, l) \sum_{r=1}^{n} C_{r,l},$$

where

$$C_{r,l} = P(0 < Z_{r-1} < l).$$

We shall now show that if $\sum_j j^2 p_j < \infty$, then

$$\limsup_n (\log n)^{-1} \sum_{l=1}^{\infty} l^{-1} B_{n,l} = 0. \tag{1}$$

To do this we need the following facts proved in [1]. If $1 < \sum_1^\infty j^2 p_j < \infty$, then for $i \ge 1$ as $r \to \infty$,

$$r P_i(0 < Z_r) \to i\lambda \tag{2}$$

and

$$\sup_{x>0} |P_i(0 < Z_n < nx | Z_n > 0) - \phi(x)| \to 0, \tag{3}$$

where $\phi(x) = 1 - e^{-\lambda x}$, $\lambda = 2\sigma^{-2}$, $\sigma^2$ being the variance of $\{p_j\}$. Since $C_{r,l} \le$

$P(0 < Z_{r-1})$, (2) implies

$$\sup_{\substack{n\geq 2\\ l\geq 1}} (\log n)^{-1} \sum_{r=1}^{n} C_{r,l} < \infty. \tag{4}$$

Next

$$\sum_{r=1}^{n} C_{r,l} \leq Kl + \sum_{r=Kl+1}^{n} C_r \left| P(0 < Z_{r-1} < l | Z_{r-1} > 0) - \phi\left(l(r-1)^{-1}\right)\right| + \phi(K^{-1}) \sum_{r=Kl+1}^{n} C_r,$$

where $C_r = P(Z_{r-1} > 0)$.

Using (2) and (3) and the fact that $\phi(x) \to 0$ as $x \to 0$, we get by choosing $K$ first and then letting $n \to \infty$ that for each fixed $l$,

$$(\log n)^{-1} \sum_{r=1}^{n} C_{r,l} \to 0, \quad \text{as } n \to \infty. \tag{5}$$

We know from Proposition 2 that

$$\sum_{1} l^{-1}\psi(l,l) < \infty, \quad \text{if } \sum_{j} j^2 p_j < \infty. \tag{6}$$

Now (1) follows from (4), (5), (6) and the dominated convergence theorem. Returning now to the inequality

$$E(Z_n\colon Z_n \geq l) \leq A_{n,l} + B_{n,l} \leq \rho l P(M_n \geq l) + B_{n,l},$$

we have on adding over $l$,

$$\rho \sum_{l} P(M_n \geq l) + \sum_{l} l^{-1} B_{n,l} \geq \sum_{l} l^{-1} E(Z_n\colon Z_n \geq l),$$

yielding the inequality

$$\rho(\log n)^{-1} EM_n + (\log n)^{-1} \sum_{l} l^{-1} B_{n,l} \geq (\log n)^{-1} E\left( Z_n \sum_{1}^{Z_n} l^{-1} \right).$$

We know from (ii) in Proposition 1 that $\sum j^2 p_j < \infty$ implies

$$(\log n)^{-1} E_i Z_n \log Z_n \to i.$$

This with (1) and the above inequality yield

$$\liminf_{n} \rho(\log n)^{-1} E_i M_n \geq i.$$

Since $\rho > 1$ is arbitrary, the proof of Proposition 4 is complete. □

### 3. Extensions.

(a) *Markov branching processes.* Our Theorem 1 extends to any continuous-time Markov branching process with finite second moments. To deduce this from Theorem 1, we proceed as follows. The upper bound of Step 1 of Section 2 can be shown to hold for any continuous-time nonnegative martingale $\{Z_t\colon t \in [0, \infty)\}$ such that $EZ_t \log Z_t \to \infty$ and thus is valid for a critical Markov branching process. Also Yaglom's theorem holds in the continuous case (see [1],

Chapter 3) and so Proposition 1 extends here. Combining these two, we get $\limsup_t E_i M_t (\log t)^{-1} \le i$. Finally, for $n \le t < n+1$ we have

$$M_t \ge M_n^* = \max_{0 \le j \le n} Z_j,$$

yielding

$$\liminf_t E_i M_t (\log t)^{-1} \ge \liminf_n E_i M_n^* (\log n)^{-1} \ge i,$$

by applying Theorem 1 to $\{Z_j\colon\ j = 0, 1, 2 \dots\}$ to get the last inequality.

A special case of the above result, namely, that of a critical birth and death process has been proved by Hammerle and Schuh in [4]. They use their result to establish a special case of our Theorem 1, namely, that of a linear fractional offspring distribution by embedding it in a critical birth and death process.

(b) *Multitype Galton–Watson process.* The methods of proof of Theorem 1 carry over to the multitype case. We state the result without proof.

THEOREM 2. *Let $\{\mathbf{Z}_n\colon\ n \ge 0\}$ be a p-type positively regular critical branching process with mean matrix $M$ and finite second moments. Let $\mathbf{u}$ and $\mathbf{v}$ be right and left eigenvectors of $M$ for the eigenvalue one normalized so that $\mathbf{u} \cdot \mathbf{1} = 1$, $\mathbf{u} \cdot \mathbf{v} = 1$, where $\mathbf{1}$ is a vector with all coordinates equal to one and $\cdot$ denotes dot product. Let $M_n = \max_{0 \le j \le n} \mathbf{v} \cdot \mathbf{Z}_j$. Then $(\log n)^{-1} E_{\mathbf{i}} M_n \to \mathbf{i} \cdot \mathbf{v}$ as $n \to \infty$.*

**Acknowledgments.** The author thanks the referee for a careful reading of the earlier versions of this paper and for making several useful suggestions. Professor H. J. Schuh caught an error in the proof of Proposition 4 in an earlier version. Thanks are due him for this and for sending me a preprint of his paper [4] with Hammerle.

## REFERENCES

[1] ATHREYA, K. B. and NEY, P. (1972). *Branching Processes*. Springer, Berlin.
[2] DOOB, J. L. (1953). *Stochastic Processes*. Wiley, New York.
[3] ERDÖS, P. (1949). On a theorem of Hsu and Robbins. *Ann. Math. Statist.* **20** 286–291.
[4] HAMMERLE, K. and SCHUH, H. J. (1986). The maximum in critical Galton–Watson and birth and death processes. *J. Appl. Probab.* **23** 601–613.
[5] HSU, P. L. and ROBBINS, H. (1947). Complete convergence and the law of large numbers. *Proc. Nat. Acad. Sci. U.S.A.* **33** 25–31.
[6] KIEFER, J. and WOLFOWITZ, J. (1956). On the characteristics of the general queueing process with applications to random walk. *Ann. Math. Statist* **27** 147–161.
[7] PAKES, A. G. (1987). Remarks on the maxima of a martingale sequence with application to the simple critical branching process. *J. Appl. Probab.* **24** 768–772.
[8] SPITZER, F. (1976). *Principles of Random Walk*, 2nd ed. Springer, New York.
[9] WEINER, H. (1984). Moments of the maximum in a critical branching process. *J. Appl. Probab.* **21** 920–923.

DEPARTMENTS OF MATHEMATICS AND STATISTICS
IOWA STATE UNIVERSITY
AMES, IOWA 50011